

Early Vascular Access PICC Candidates Assessment Advantage Program™

Diagnosis		
Abscess	Craniotomy	Pancreatic cancer
Acute leukemia	Cystic fibrosis	Pancreatitis
Acute renal failure	Decubitis ulcers	Pericarditis
AIDS	Diabetes insipidus	Peritonitis
Burns	Disseminated Intravascular Coagulation	Pneumocystis pneumonia
Cardiomyopathy	Emphysema (COPD)	Post op wound infection
Cellulitis	Encephalitis	Pseudomonas infection
Cerebral aneurysm	Endocarditis	Ruptured appendix
Cerebral vascular accident	Hyperemesis	Septic arthritis
Chron's disease	Inter-cerebral hematoma	Septicemia
Cirrhosis	Liver transplant	Small and large bowel procedures—diverticulitis, intestinal obstruction
CMV	Lyme disease	Toxoplasmosis
Complicated pneumonia	Meningitis	Ulcerative Colitis
Complicated pneumonia—ventilator support	Myocarditis	Wound Infections
Congestive Heart Failure	Osteomyelitis	

Medical Conditions/Patient History		
Diabetes	Low platelet count	Multiple hospital admissions
Fluid restricted patients	Multiple blood draws	Pain control
IV drug abuse	Multiple drugs—compatible	Steroids
Limited venous access on admission	Multiple drugs—incompatible	

Drugs		
5 – FU® pH 9.2	Dopamine pH 2.5 – 5	Phenytoin pH 12
Acyclovir pH 10.5-11.6	Doxycycline pH 1.8-3.3	Potassium pH 4.0
Amphotericin—irritant	Erythromycin—irritant	PPN & TPN—Hypertonic
Ampicillin pH 8.0 -10	Ganciclovir pH 9-11	Taxol® pH 4.4-5.6
Bactrim® pH 10	Nafcillin—Hypertonic	Vancomycin pH 2.4
Ciprofloxacin pH 3.3 - 4.6	Pentamidine pH 4.1 – 5.4	